

YOUR CARE TEAM DISCUSSION GUIDE



It can be hard to start conversations about your treatment with your care team. This guide can help you keep track of questions you want to ask and topics to discuss so you can feel confident going into your appointments.

What is ZIIHERA?

ZIIHERA (zanidatamab-hrii) 50 mg/mL is a prescription medicine used to treat adults with gastroesophageal cancer (GEA) including gastric cancer, esophageal cancer, and gastroesophageal junction (GEJ) cancer. ZIIHERA is used in combination with fluoropyrimidine- and platinum-containing chemotherapy with or without tislelizumab-jsgr as the first treatment when your GEA:

- cannot be removed by surgery or has spread to other parts of your body (metastatic); **and**
- your tumor tests positive for human epidermal growth factor receptor 2 (HER2 IHC 3+ or IHC 2+/ISH+).

Your healthcare provider will perform a test to make sure ZIIHERA is right for you.

It is not known if ZIIHERA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

ZIIHERA can cause serious side effects, including:

- **Diarrhea.** Diarrhea during treatment with ZIIHERA can be severe, life threatening, and can lead to death. Diarrhea may cause loss of fluids and body salts (electrolytes). To help prevent or reduce diarrhea during treatment with ZIIHERA, your healthcare provider:
 - will tell you to drink more fluids.
 - start you on an antidiarrheal medicine such as loperamide or other treatment, as needed.
 - may temporarily stop, reduce your dose or permanently stop treatment with ZIIHERA, or other medicines you take.

If you get new or worsening diarrhea, tell your healthcare provider right away and take antidiarrheal medicine as prescribed. Follow all of your healthcare providers instructions to help with your diarrhea.

- **Harm to your unborn baby.** Taking ZIIHERA during pregnancy or within 4 months before pregnancy can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZIIHERA.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ZIIHERA.
- Use effective birth control (contraception) during treatment with ZIIHERA and for 4 months after the last dose.

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.


ZIIHERA[®]
(zanidatamab-hrii)
50mg/mL Injection for IV

CONVERSATIONS WITH YOUR CARE TEAM BEGIN BEFORE YOUR APPOINTMENT



YOUR CARE TEAM IS HERE TO HELP YOU WITH YOUR ZIIHERA (zanidatamab-hrii) JOURNEY

It can be hard to know where to begin when discussing your treatment with your care team. This guide can help you keep track of questions you want to ask. It can also give you ideas of what you want to discuss.

Working closely with your care team can help you navigate your treatment journey. Your care team is an excellent resource for answering your questions and connecting you with additional support and information.

Now that you've been prescribed ZIIHERA as part of your treatment plan, your care team can help you understand what your treatment will look like.



QUESTIONS YOU CAN ASK YOUR CARE TEAM ABOUT YOUR DIAGNOSIS:

☐ Where in my body did my cancer start? _____

☐ What is biomarker testing and what does it mean that my cancer is HER2-positive? _____

Space for notes and other questions you may have for your care team:

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including **BOXED WARNINGS** and **Medication Guide**, and discuss with your doctor.


ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

PREPARING FOR YOUR APPOINTMENTS CAN HELP MAKE THEM LESS STRESSFUL



QUESTIONS TO ASK YOUR CARE TEAM ABOUT YOUR TREATMENT:

☐ Why did you choose this treatment for me? _____

☐ How will we know if this treatment is working for me? _____

☐ What side effects can I expect from this treatment? _____

☐ Is there anything I can do ahead of time to be prepared for potential side effects? _____

☐ Who should I call if I am having problems from my treatment, and how do I know if I need to call them? _____

Space for notes and other questions you may have for your care team:

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.


ZIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

SHARE WHAT MATTERS TO YOU AS PART OF YOUR TREATMENT JOURNEY

Let your care team know what your goals are and what matters to you. The more they know, the better they can help. You can talk to them about what you want your life to look like while you're taking ZIIHERA (zanidatamab-hrii)-based treatment.

WHAT MOTIVATES ME:

 Visiting _____

 Spending time with _____

 Celebrating _____

 Learning _____

 Doing _____

What are other things that are important to you?

**THINKING ABOUT WHAT YOU WANT TO TALK ABOUT AHEAD OF TIME CAN
HELP YOU HAVE CLEARER CONVERSATIONS WITH YOUR CARE TEAM.**

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.


ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

Other questions you may have for your care team:

Notes and other things you want to remember:



LEARN MORE ABOUT ZIIHERA

Scan the QR code to visit ZIIHERA.com
and download your Treatment Brochure.

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.


ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

APPROVED USE AND IMPORTANT SAFETY INFORMATION

What is ZIIHERA?

ZIIHERA (zanidatamab-hrii) 50 mg/mL is a prescription medicine used to treat adults with gastroesophageal cancer (GEA) including gastric cancer, esophageal cancer, and gastroesophageal junction (GEJ) cancer. ZIIHERA is used in combination with fluoropyrimidine- and platinum-containing chemotherapy with or without tislelizumab-jsgr as the first treatment when your GEA:

- cannot be removed by surgery or has spread to other parts of your body (metastatic); **and**
- your tumor tests positive for human epidermal growth factor receptor 2 (HER2 IHC 3+ or IHC 2+/ISH+).

Your healthcare provider will perform a test to make sure ZIIHERA is right for you.

It is not known if ZIIHERA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

ZIIHERA can cause serious side effects, including:

- **Diarrhea.** Diarrhea during treatment with ZIIHERA can be severe, life threatening, and can lead to death. Diarrhea may cause loss of fluids and body salts (electrolytes). To help prevent or reduce diarrhea during treatment with ZIIHERA, your healthcare provider:
 - will tell you to drink more fluids.
 - start you on an antidiarrheal medicine such as loperamide or other treatment, as needed.
 - may temporarily stop, reduce your dose or permanently stop treatment with ZIIHERA, or other medicines you take.

If you get new or worsening diarrhea, tell your healthcare provider right away and take antidiarrheal medicine as prescribed. Follow all of your healthcare providers instructions to help with your diarrhea.

- **Harm to your unborn baby.** Taking ZIIHERA during pregnancy or within 4 months before pregnancy can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZIIHERA.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ZIIHERA.
- Use effective birth control (contraception) during treatment with ZIIHERA and for 4 months after the last dose.
- **Heart problems** that may affect how well your heart pumps blood. Your healthcare provider will check your heart function before and during treatment with ZIIHERA. Tell your healthcare provider right away if you get any of the following signs and symptoms of a heart problem:
 - new or worsening shortness of breath
 - irregular heartbeat
 - feeling more tired than usual
 - sudden weight gain
 - swelling of your ankles or feet
 - dizziness or feeling light-headed
 - loss of consciousness

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.



APPROVED USE AND IMPORTANT SAFETY INFORMATION (CONT'D)

- **Infusion-related reactions.** Your healthcare provider will check for side effects during your infusions. Tell your healthcare provider right away if you get any of the following symptoms during or after your infusions of ZIIHERA:

- shortness of breath or trouble breathing
- fever
- chills
- rash
- flushing
- nausea or vomiting
- dizziness or feeling light-headed
- chest discomfort

Before receiving ZIIHERA, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had any heart problems.
- are breastfeeding or plan to breastfeed. It is not known if ZIIHERA passes into your breastmilk. Talk to your healthcare provider about the best way to feed your baby if you receive treatment with ZIIHERA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Other common side effects when ZIIHERA is given in combination with chemotherapy and tislelizumab-jsgr include:

- diarrhea
- nausea
- decreased appetite
- low levels of blood potassium
- vomiting
- feeling tired
- numbness, pain, tingling, or burning in your hands or feet
- rash
- infusion-related reactions

Other common side effects when ZIIHERA is given in combination with chemotherapy include:

- diarrhea
- nausea
- vomiting
- numbness, pain, tingling, or burning in your hands or feet
- decreased appetite
- feeling tired
- low levels of blood potassium
- infusion-related reactions
- rash

These are not all the possible side effects of ZIIHERA. Call your doctor for medical advice about side effects.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. You may report side effects to FDA at 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.


ZIIHERA[®]
(zanidatamab-hrii)
50mg/mL Injection for IV



ZIIHERA IS

HERE FOR WHAT'S NEXT



ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

TIPS FOR GETTING THE MOST OUT OF YOUR APPOINTMENTS:

- ✓ Keep track of how you feel between visits
- ✓ Write down questions you have as you think of them to help you keep track of what you want to ask your care team
- ✓ Be open with your care team so they can better manage your treatment
- ✓ Bring someone to help support you during your appointments. They can help you take notes and ask questions

What is ZIIHERA?

ZIIHERA (zanidatamab-hrii) 50 mg/mL is a prescription medicine used to treat adults with gastroesophageal cancer (GEA) including gastric cancer, esophageal cancer, and gastroesophageal junction (GEJ) cancer. ZIIHERA is used in combination with fluoropyrimidine- and platinum-containing chemotherapy with or without tislelizumab-jsgr as the first treatment when your GEA:

- cannot be removed by surgery or has spread to other parts of your body (metastatic); **and**
- your tumor tests positive for human epidermal growth factor receptor 2 (HER2 IHC 3+ or IHC 2+/ISH+).

Your healthcare provider will perform a test to make sure ZIIHERA is right for you.

It is not known if ZIIHERA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

ZIIHERA can cause serious side effects, including:

- **Diarrhea.** Diarrhea during treatment with ZIIHERA can be severe, life threatening, and can lead to death. Diarrhea may cause loss of fluids and body salts (electrolytes). To help prevent or reduce diarrhea during treatment with ZIIHERA, your healthcare provider:
 - will tell you to drink more fluids.
 - start you on an antidiarrheal medicine such as loperamide or other treatment, as needed.
 - may temporarily stop, reduce your dose or permanently stop treatment with ZIIHERA, or other medicines you take.

If you get new or worsening diarrhea, tell your healthcare provider right away and take antidiarrheal medicine as prescribed. Follow all of your healthcare providers instructions to help with your diarrhea.

- **Harm to your unborn baby.** Taking ZIIHERA during pregnancy or within 4 months before pregnancy can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZIIHERA.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ZIIHERA.
- Use effective birth control (contraception) during treatment with ZIIHERA and for 4 months after the last dose.

Please see additional Important Safety Information throughout, and accompanying full Prescribing Information, including BOXED WARNINGS and Medication Guide, and discuss with your doctor.



Jazz Pharmaceuticals.

©2026 Jazz Pharmaceuticals plc or its subsidiaries US-ZAN-2600042