

WHAT TO EXPECT FROM TREATMENT WITH ZIIHERA

WHAT TO EXPECT GUIDE

A guide to help you prepare for treatment.

AI-generated synthetic performer; not a real patient.

What is ZIIHERA?

ZIIHERA (zanidatamab-hrii) 50 mg/mL is a prescription medicine used to treat adults with gastroesophageal cancer (GEA) including gastric cancer, esophageal cancer, and gastroesophageal junction (GEJ) cancer. ZIIHERA is used in combination with fluoropyrimidine- and platinum-containing chemotherapy with or without tislelizumab-jsgx as the first treatment when your GEA:

- cannot be removed by surgery or has spread to other parts of your body (metastatic); **and**
- your tumor tests positive for human epidermal growth factor receptor 2 (HER2 IHC 3+ or IHC 2+/ISH+).

Your healthcare provider will perform a test to make sure ZIIHERA is right for you.

It is not known if ZIIHERA is safe and effective in children.

IMPORTANT SAFETY INFORMATION

ZIIHERA can cause serious side effects, including:

- **Diarrhea.** Diarrhea during treatment with ZIIHERA can be severe, life threatening, and can lead to death. Diarrhea may cause loss of fluids and body salts (electrolytes). To help prevent or reduce diarrhea during treatment with ZIIHERA, your healthcare provider:
 - will tell you to drink more fluids.
 - start you on an antidiarrheal medicine such as loperamide or other treatment, as needed.
 - may temporarily stop, reduce your dose or permanently stop treatment with ZIIHERA, or other medicines you take.

Please see Important Safety Information throughout and full Prescribing Information, including BOXED WARNINGS and Medication Guide.




ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

STARTING ZIIHERA



Starting a new treatment can feel easier when you know what is ahead

You don't have to navigate this alone. Your care team is here to help support you at every step. This guide is here to help you take charge of your care and help you feel empowered for what's to come.



Throughout this brochure, you'll learn:

- **How ZIIHERA is given**
- **How often you may receive treatment**
- **Which side effects may occur and how they may be managed**

IMPORTANT SAFETY INFORMATION

ZIIHERA can cause serious side effects, including (cont'd):

If you get new or worsening diarrhea, tell your healthcare provider right away and take antidiarrheal medicine as prescribed. Follow all of your healthcare providers instructions to help with your diarrhea.

- **Harm to your unborn baby.** Taking ZIIHERA during pregnancy or within 4 months before pregnancy can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZIIHERA.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ZIIHERA.
- Use effective birth control (contraception) during treatment with ZIIHERA and for 4 months after the last dose.

Please see Important Safety Information throughout and full Prescribing Information, including **BOXED WARNINGS and Medication Guide.**

TIPS FOR TREATMENT



Starting treatment is a big step. As you begin your journey, **there are simple things you can do each day to feel more like yourself.** Remember, small steps add up to make a real difference.

Eating well

- Try to eat small, frequent meals throughout the day. Soft, moist, high-protein foods like eggs and fish help keep your strength up

Managing swallowing

- Blended or pureed foods like smoothies, soups, and soft-cooked options can make eating more comfortable

Staying hydrated

- Staying hydrated is important. Sip fluids throughout the day

Daily planning

- Prep meals and write down questions for your doctor on days when you are feeling well. Lean on friends and family for help when you need it

Emotional health

- It is okay to feel scared or overwhelmed. Talking to someone you trust, or connecting with a support group, can help you feel less alone

Staying active

- Exercise can help you feel better and keep your energy up. Ask your doctor what level of activity is right for you

IMPORTANT SAFETY INFORMATION (CONT'D)

ZIIHERA can cause serious side effects, including (cont'd):

- **Heart problems** that may affect how well your heart pumps blood. Your healthcare provider will check your heart function before and during treatment with ZIIHERA. Tell your healthcare provider right away if you get any of the following signs and symptoms of a heart problem:
 - new or worsening shortness of breath
 - feeling more tired than usual
 - swelling of your ankles or feet
 - loss of consciousness
 - irregular heartbeat
 - sudden weight gain
 - dizziness or feeling light-headed

Please see Important Safety Information throughout and full [Prescribing Information](#), including **BOXED WARNINGS** and [Medication Guide](#).

BEFORE YOUR INFUSION


ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV



ZIIHERA is given as an intravenous (IV) infusion, which means the medicine is delivered directly into your bloodstream.

You will receive ZIIHERA **every 2 or 3 weeks** in a clinic or an infusion center. Your doctor will decide what schedule is right for you.

You will receive chemotherapy along with ZIIHERA during the same visit. **Your doctor may also decide to add immunotherapy to your treatment plan.** Your care team will decide what and how many treatments you need.

- **Immunotherapy:** a treatment that may help activate the immune system so your body can fight cancer

You will receive some medicines 30 to 60 minutes before each treatment to reduce the risk of infusion-related reactions or make them less severe. Your care team will also direct you to take an over-the-counter medicine called loperamide (Imodium[®]*) to help manage diarrhea.

Have questions before your treatment? Our Care Team Discussion Guide can help you track your questions and thoughts. To download one, scan the QR code or visit ZIIHERA.com/resources



*Imodium is a registered trademark of Kenvue Brands LLC.

IMPORTANT SAFETY INFORMATION (CONT'D)

ZIIHERA can cause serious side effects, including (cont'd):

- **Infusion-related reactions.** Your healthcare provider will check for side effects during your infusions. Tell your healthcare provider right away if you get any of the following symptoms during or after your infusions of ZIIHERA:
 - shortness of breath or trouble breathing
 - fever
 - chills
 - rash
 - flushing
 - nausea or vomiting
 - dizziness or feeling light-headed
 - chest discomfort

Please see Important Safety Information throughout and full Prescribing Information, including **BOXED WARNINGS and Medication Guide.**

DURING YOUR INFUSION


ZIIHERA[®]
(zanidatamab-hrii)
50mg/ml Injection for IV

- Plan for your ZIIHERA infusions to take some time. Your first treatment will likely take the longest
- You may want to bring a book, download a podcast, or plan to watch a TV show during your infusions



Your care team will check if you are having any side effects during your infusion. You should tell them right away if you have any of these symptoms at any time during or after your infusions:

- shortness of breath or trouble breathing
- fever
- chills
- rash
- flushing
- nausea or vomiting
- dizziness or feeling lightheaded
- chest discomfort

As you continue treatment, your care team will monitor for any side effects you may have. If you miss a dose, call your healthcare provider as soon as possible to reschedule your appointment.


IMPORTANT SAFETY INFORMATION (CONT'D)

Before receiving ZIIHERA, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had any heart problems.
- are breastfeeding or plan to breastfeed. It is not known if ZIIHERA passes into your breastmilk. Talk to your healthcare provider about the best way to feed your baby if you receive treatment with ZIIHERA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout and full Prescribing Information, including BOXED WARNINGS and Medication Guide.



SIDE EFFECTS



KNOWING WHAT TO EXPECT HELPS YOU PREPARE FOR YOUR TREATMENT

ZIIHERA can cause serious side effects, including:

- **Diarrhea.** Diarrhea during treatment with ZIIHERA can be severe, life threatening, and can lead to death. Diarrhea may cause loss of fluids and body salts (electrolytes). Follow all of your care team's instructions to help with your diarrhea.
- **Harm to an unborn baby.** Taking ZIIHERA during pregnancy or within 4 months before pregnancy can harm your unborn baby.
- **Heart problems** that may affect how well your heart pumps blood.
- **Infusion-related reactions**

More severe or worse side effects occurred more often in people who were 65 years or older who received ZIIHERA + tislelizumab (immunotherapy) + chemotherapy (87%) compared to people younger than 65 years (80%).

Other common side effects when ZIIHERA is given in combination with chemotherapy and tislelizumab include:

- diarrhea
- nausea
- decreased appetite
- low levels of blood potassium
- vomiting
- feeling tired
- numbness, pain, tingling, or burning in your hands or feet
- rash
- infusion-related reactions

Other common side effects when ZIIHERA is given in combination with chemotherapy include:

- diarrhea
- nausea
- vomiting
- numbness, pain, tingling, or burning in your hands or feet
- decreased appetite
- feeling tired
- low levels of blood potassium
- infusion-related reactions
- rash

Tell your care team if you have any side effect that bothers you or that does not go away.

- Your care team may slow down, pause, or stop your treatment with ZIIHERA if you have certain side effects. They will work with you to help manage side effects and treat them when they do happen.

Please see Important Safety Information throughout and full [Prescribing Information](#), including **BOXED WARNINGS and [Medication Guide](#).**

SIDE EFFECTS (CONT'D)



Managing diarrhea:

With ZIIHERA treatment plans, **about 4 out of 5 people** experienced diarrhea. Your care team expects this and can help you manage it.

- For most people, diarrhea happened during the first round of treatment and got better within 3 weeks. Diarrhea may happen again in later rounds of treatment as well
- If you get new or worsening diarrhea, notify your care team right away

To help manage diarrhea, your care team will give you an over-the-counter medicine called **loperamide (Imodium[®]*)**.

- Take 4 mg twice a day, starting before you receive ZIIHERA on the day of your first treatment for at least the first 7 days. Your doctor may ask you to keep taking it after that, if needed



Drinking extra fluids is important since diarrhea can cause your body to lose water. **Don't try to push through diarrhea on your own.** Call your care team right away if you experience frequent or loose stools or if anything feels unmanageable. They are there to help.



Be sure to track any side effects you may have in your **ZIIHERA Journey Tracker**

ZIIHERA.com/resources

*Imodium is a registered trademark of Kenvue Brands LLC.

IMPORTANT SAFETY INFORMATION (CONT'D)

Other common side effects when ZIIHERA is given in combination with chemotherapy and tislelizumab-jsgr include:

- | | |
|---------------------------------|--|
| ○ diarrhea | ○ feeling tired |
| ○ nausea | ○ numbness, pain, tingling, or burning in your hands or feet |
| ○ decreased appetite | ○ rash |
| ○ low levels of blood potassium | ○ infusion-related reactions |
| ○ vomiting | |

Please see Important Safety Information throughout and full Prescribing Information, including BOXED WARNINGS and Medication Guide.

IMPORTANT SAFETY INFORMATION



IMPORTANT SAFETY INFORMATION

ZIIHERA can cause serious side effects, including:

- **Diarrhea.** Diarrhea during treatment with ZIIHERA can be severe, life threatening, and can lead to death. Diarrhea may cause loss of fluids and body salts (electrolytes). To help prevent or reduce diarrhea during treatment with ZIIHERA, your healthcare provider:
 - will tell you to drink more fluids.
 - start you on an antidiarrheal medicine such as loperamide or other treatment, as needed.
 - may temporarily stop, reduce your dose or permanently stop treatment with ZIIHERA, or other medicines you take.

If you get new or worsening diarrhea, tell your healthcare provider right away and take antidiarrheal medicine as prescribed. Follow all of your healthcare providers instructions to help with your diarrhea.

- **Harm to your unborn baby.** Taking ZIIHERA during pregnancy or within 4 months before pregnancy can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with ZIIHERA.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with ZIIHERA.
- Use effective birth control (contraception) during treatment with ZIIHERA and for 4 months after the last dose.
- **Heart problems** that may affect how well your heart pumps blood. Your healthcare provider will check your heart function before and during treatment with ZIIHERA. Tell your healthcare provider right away if you get any of the following signs and symptoms of a heart problem:
 - new or worsening shortness of breath
 - irregular heartbeat
 - feeling more tired than usual
 - sudden weight gain
 - swelling of your ankles or feet
 - dizziness or feeling light-headed
 - loss of consciousness
- **Infusion-related reactions.** Your healthcare provider will check for side effects during your infusions.

Please see Important Safety Information throughout and full [Prescribing Information](#), including **BOXED WARNINGS** and [Medication Guide](#).

IMPORTANT SAFETY INFORMATION (CONT'D)



Tell your healthcare provider right away if you get any of the following symptoms during or after your infusions of ZIIHERA:

- shortness of breath or trouble breathing
- fever
- chills
- rash
- flushing
- nausea or vomiting
- dizziness or feeling light-headed
- chest discomfort

Before receiving ZIIHERA, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had any heart problems.
- are breastfeeding or plan to breastfeed. It is not known if ZIIHERA passes into your breastmilk. Talk to your healthcare provider about the best way to feed your baby if you receive treatment with ZIIHERA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Other common side effects when ZIIHERA is given in combination with chemotherapy and tislelizumab-jsgr include:

- diarrhea
- nausea
- decreased appetite
- low levels of blood potassium
- vomiting
- feeling tired
- numbness, pain, tingling, or burning in your hands or feet
- rash
- infusion-related reactions

Other common side effects when ZIIHERA is given in combination with chemotherapy include:

- diarrhea
- nausea
- vomiting
- numbness, pain, tingling, or burning in your hands or feet
- decreased appetite
- feeling tired
- low levels of blood potassium
- infusion-related reactions
- rash

These are not all the possible side effects of ZIIHERA. Call your doctor for medical advice about side effects.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see Important Safety Information throughout and full Prescribing Information, including **BOXED WARNINGS and Medication Guide.**

NOTES



ZIHERA®
(zanidatamab-hrii)
50mg/ml Injection for IV

Space for you to collect your thoughts, questions, and anything else on your mind:



Please see Important Safety Information throughout and full [Prescribing Information](#), including **BOXED WARNINGS and [Medication Guide](#).**



ZIIHERA CAN SUPPORT YOU ON YOUR NEXT STEP

Talk to your doctor and
visit **ZIIHERA.com**
to discover more.



AI-generated synthetic performer; not a real patient.

IMPORTANT SAFETY INFORMATION (CONT'D)

Other common side effects when ZIIHERA is given in combination with chemotherapy include:

- diarrhea
- nausea
- vomiting
- numbness, pain, tingling, or burning in your hands or feet
- decreased appetite
- feeling tired
- low levels of blood potassium
- infusion-related reactions
- rash

These are not all the possible side effects of ZIIHERA. Call your doctor for medical advice about side effects.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. You may report side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see Important Safety Information throughout and full Prescribing Information, including BOXED WARNINGS and Medication Guide.